

# Florida, 2013

## Primary and secondary syphilis

- CASES: Florida reported 1,513 cases in 2013.
- RATES: Florida ranked 4 among 50 states reporting at least one case of P&S syphilis with 7.8 cases per 100,000 population compared to the U.S. rate of 5.5.

## Rates of primary and secondary syphilis, by sex

- In Florida, the rate among males was 14.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.4 compared to the U.S. female rate of 0.9.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Florida, the race/ethnicity adjusted rates per 100,000 population were 4.3 among whites, 18.5 among blacks, 8.4 among Hispanics, 2.2 among Asians, 8.3 among Native Hawaiian/Other Pacific Islanders, and 4.0 among American Indians/Alaska Natives.
- The rate among blacks was 4.4 times that of whites.
- The rate among Hispanics was 2 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 2 times that of whites.


## Congenital syphilis

- CASES: Florida reported 37 cases in 2013.
- RATES: Florida ranked 5 among 25 states reporting at least one case of congenital syphilis with a rate of 17.2 cases per 100,000 live births compared to the U.S. rate of 8.7.

## Syphilis: P&S Rates and Congenital Cases, Florida, 2004–2013



## Total P&S Syphilis Rates by County, Florida, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2